

Alert	1:10,000 (1 mg/10 mL) ampoule is the preferred preparation for adrenaline infusion.												
Indication	Treatment of hypotensive shock with or without myocardial dysfunction.												
Action	<p>Catecholamine with alpha and beta adrenergic actions. Haemodynamic effects are dose dependent:</p> <ul style="list-style-type: none"> At low doses of 0.01–0.1 microgram/kg/minute primarily stimulates cardiac and vascular beta 1- and beta 2-adrenoreceptors leading to increased inotropy, chronotropy, conduction velocity and peripheral vasodilation. At doses greater than 0.1 microgram/kg/minute adrenaline also stimulates vascular and cardiac alpha 1-receptors causing vasoconstriction and increased inotropy. The net effects are increases in blood pressure and systemic blood flow caused by the drug-induced increases in systemic vascular resistance (SVR) and cardiac output.¹ 												
Drug Type	Inotropic vasopressor.												
Trade Name	Aspen Adrenaline 1: 10,000 Adrenaline Acid Tartrate injection; Adrenaline 1:1,000 Adrenalin Acid Tartrate injection.												
Presentation	1 mg/10 mL or 1:10,000 ampoule [100 microgram/mL] 1 mg/mL or 1:1,000 ampoule [1000 microgram/mL]												
Dosage / Interval	Low dose: 0.05–0.1 microgram/kg/minute High dose: 0.1–1 microgram/kg/minute												
Route	Continuous IV infusion.												
Preparation/Dilution	<p>Preparation using 1:10,000 (1 mg/10 mL) ampoule</p> <p>LOW CONCENTRATION IV infusion</p> <table border="1"> <thead> <tr> <th>Infusion dose</th> <th>Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.05 microgram/kg/minute</td> <td>150 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 150 microgram/kg [1.5 mL/kg] of 1:10,000 adrenaline and add glucose 5%, glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 3 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.05 microgram/kg/minute.</p> <p>HIGH CONCENTRATION IV infusion</p> <table border="1"> <thead> <tr> <th>Infusion dose</th> <th>Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.2 microgram/kg/minute</td> <td>600 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 600 microgram/kg [6 mL/kg] of 1:10,000 adrenaline and add glucose 5%, glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 12 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.2 microgram/kg/minute.</p> <p>For infants requiring fluid restriction consider: VERY HIGH CONCENTRATION IV infusion*</p> <table border="1"> <thead> <tr> <th>Infusion dose</th> <th>Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.4 microgram/kg/minute</td> <td>1200 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 1200 microgram/kg [12 mL/kg] of 1:10,000 adrenaline and add glucose 5% ONLY to make a final volume of 50 mL with a concentration of 24 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.4 microgram/kg/minute.</p> <p>*Stability data only available for 5% glucose for very high concentration.</p>	Infusion dose	Prescribed amount	1 mL/hour = 0.05 microgram/kg/minute	150 microgram/kg adrenaline and make up to 50 mL	Infusion dose	Prescribed amount	1 mL/hour = 0.2 microgram/kg/minute	600 microgram/kg adrenaline and make up to 50 mL	Infusion dose	Prescribed amount	1 mL/hour = 0.4 microgram/kg/minute	1200 microgram/kg adrenaline and make up to 50 mL
Infusion dose	Prescribed amount												
1 mL/hour = 0.05 microgram/kg/minute	150 microgram/kg adrenaline and make up to 50 mL												
Infusion dose	Prescribed amount												
1 mL/hour = 0.2 microgram/kg/minute	600 microgram/kg adrenaline and make up to 50 mL												
Infusion dose	Prescribed amount												
1 mL/hour = 0.4 microgram/kg/minute	1200 microgram/kg adrenaline and make up to 50 mL												

	<p>Preparation using 1:1,000 (1 mg/mL) ampoule – Occasionally used for infants >4 kg. : 1:1000 (1 mg/mL) ampoule is generally not kept in the NICUs</p> <p>LOW CONCENTRATION IV infusion</p> <table border="1"> <thead> <tr> <th>Infusion dose</th> <th>Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.05 microgram/kg/minute</td> <td>150 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 150 microgram/kg [0.15 mL/kg] of 1:1000 adrenaline and add glucose 5%, glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 3 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.05 microgram/kg/minute.</p> <p>HIGH CONCENTRATION IV infusion</p> <table border="1"> <thead> <tr> <th>Infusion dose</th> <th>Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.2 microgram/kg/minute</td> <td>600 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 600 microgram/kg [0.6 mL/kg] of 1:1000 adrenaline and add glucose 5%, glucose 10% or sodium chloride 0.9% to make a final volume of 50 mL with a concentration of 12 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.2 microgram/kg/minute.</p> <p>For infants requiring fluid restriction consider: VERY HIGH CONCENTRATION IV infusion*</p> <table border="1"> <thead> <tr> <th>Infusion dose</th> <th>Prescribed amount</th> </tr> </thead> <tbody> <tr> <td>1 mL/hour = 0.4 microgram/kg/minute</td> <td>1200 microgram/kg adrenaline and make up to 50 mL</td> </tr> </tbody> </table> <p>Draw up 1200 microgram/kg [1.2 mL/kg] of 1:1000 adrenaline and add glucose 5% ONLY to make a final volume of 50 mL with a concentration of 24 microgram/kg/mL. Infusing at a rate of 1 mL/hour = 0.4 microgram/kg/minute. *Stability data only available for 5% glucose for very high concentration.</p>	Infusion dose	Prescribed amount	1 mL/hour = 0.05 microgram/kg/minute	150 microgram/kg adrenaline and make up to 50 mL	Infusion dose	Prescribed amount	1 mL/hour = 0.2 microgram/kg/minute	600 microgram/kg adrenaline and make up to 50 mL	Infusion dose	Prescribed amount	1 mL/hour = 0.4 microgram/kg/minute	1200 microgram/kg adrenaline and make up to 50 mL
Infusion dose	Prescribed amount												
1 mL/hour = 0.05 microgram/kg/minute	150 microgram/kg adrenaline and make up to 50 mL												
Infusion dose	Prescribed amount												
1 mL/hour = 0.2 microgram/kg/minute	600 microgram/kg adrenaline and make up to 50 mL												
Infusion dose	Prescribed amount												
1 mL/hour = 0.4 microgram/kg/minute	1200 microgram/kg adrenaline and make up to 50 mL												
Administration	Continuous intravenous infusion via a central line. Use with caution via a peripheral line.												
Monitoring	Continuous heart rate, ECG and blood pressure monitoring preferable. Assess urine output and peripheral perfusion frequently. Observe IV site closely for blanching and extravasation.												
Contraindications	Arrhythmia and tachyarrhythmia. Cardiovascular disease resulting in arterial narrowing including cerebrovascular disease, coronary artery disease and digital ischaemia. Pheochromocytoma. Thyrotoxicosis. Glaucoma. Known hypersensitivity to sympathomimetic amines.												
Precautions	Ensure adequate circulating blood volume prior to commencement. Adrenaline is a potent chronotrope and vasopressor – may cause excessive tachycardia, severe hypertension and ventricular arrhythmias. Adrenaline may cause lactic acidosis and hyperglycaemia.												
Drug Interactions	Hypotension may be observed with concurrent use of vasodilators such as glyceryl trinitrate, nitroprusside and calcium channel blockers. Concurrent use of digitalis glycosides may increase the risk of cardiac arrhythmias. Concurrent use of IV phenytoin with adrenaline may result in dose dependent, sudden hypotension and bradycardia.												
Adverse Reactions	Tachycardia and arrhythmia. Systemic hypertension especially at higher doses. May cause hypokalaemia. Tissue necrosis at infusion site with extravasation. Digital ischaemia.												
Compatibility	Fluids: Glucose 5%, glucose 10%, Hartmann's, sodium chloride 0.9%. Stability data only available for 5% glucose for very high concentration.												

	Y-site: Amino acid solutions. Amiodarone, anidulafungin, atracurium, bivalirudin, caspofungin, cisatracurium, dexmedetomidine, dobutamine, dopamine, ethanol, fentanyl, glyceryl trinitrate, heparin sodium, milrinone, morphine sulfate, pancuronium, potassium chloride, ranitidine, remifentanyl, sodium nitroprusside, tigecycline, tirofiban, vecuronium.
Incompatibility	<p>Fluids: Sodium bicarbonate.</p> <p>Y-site: Aciclovir, aminophylline, ampicillin, atropine, azathioprine, calcium chloride, calcium gluconate, cefalotin, chloramphenicol, digoxin, ergometrine, ganciclovir, hyaluronidase, hydrocortisone sodium succinate, indomethacin, phenobarbitone sodium, sodium bicarbonate, thiopentone, vancomycin.</p> <p>No information: Adrenaline HCL is compatible with noradrenaline bitartrate but no stability data is available for Adrenaline acid tartrate and noradrenaline bitartrate</p>
Stability	<p>Ampoule: Store below 30°C. Protect from light.</p> <p>Diluted solution: Stable for 24 hours below 25°C.</p>
Storage	Ampoule: Store below 25°C. Protect from light. Discard remainder after use.
Special Comments	<p>Ensure adrenaline has a "dedicated" line to avoid accidental bolus. Do not use as a side line with maintenance fluids.</p> <p>Discard if exhibiting colour change.</p>
Evidence summary	Refer to full version.
References	Refer to full version.

Original version Date: 31/03/2016	Author: NMF Consensus Group
Current Version number: 1.2	Current Version Date: 30/11/2017
Risk Rating: Medium	Due for Review: 30/11/2020
Approval by: JHCH CQ&PCC	Approval Date: 27/02/2018

Authors Contribution

Original author	David Osborn
Evidence Review	David Osborn
Content review of the current version	Srinivas Bolisetty, David Osborn, Eszter Jozsa, Nilkant Phad, Cindy Chen
Nursing Review	Eszter Jozsa
Pharmacy Review	Jing Xiao, Mariella De Rosa, Ushma Trivedi, Cindy Chen
Final content and editing review of the original	Ian Whyte
Electronic version	Mariella De Rosa, Cindy Chen, Ian Callander
Facilitator	Srinivas Bolisetty